

SERIAL NO.








10/783,095

PEAKMAN, Mark

TC/A.U.

1653

[illegible][illegible]

	<p>Congia, M., et al; "T cell epitopes of insulin defined in HLA-DR4 transgenic mice are derived from preproinsulin and proinsulin"; <i>Proc. Natl. Acad. Sci. USA</i>; Vol. 95, No. 7; pp. 3833-3838, 31 March 1998; XP-002204284.</p>
	<p>Geluk, A., et al; "HLA-DR Binding Analysis of Peptides from Islet Antigens in IDDM"; <i>Diabetes</i>; Vol. 47, No. 10; pp. 1594-1601, October 1998; XP-002323217.</p>
	<p>Narendran, P., et al; "Humoral and cellular immune responses to proinsulin in adults with newly diagnosed type 1 diabetes"; <i>Diabetes/Metabolism Research and Reviews</i>; Vol. 19, No. 1; pp. 52-59; 28 October 2003; XP-002323390.</p>
	<p>Liberman, S.M., et al; "A comprehensive guide to antibody and T-cell responses in type 1 diabetes"; <i>Tissue Antigens</i>; Vol. 62, No. 5; pp. 359-377; November 2003; XP-002323391.</p>
	<p>Meierhoff, G., et al; "Cytokine detection by ELISPOT: relevance for immunological studies in type 1 diabetes"; <i>Diabetes/Metabolism Research and Reviews</i>; Vol. 18, No. 5, pp. 367-380; October 2002; XP-002323218.</p>
	<p>Peakman, M., et al; "Naturally processed and presented epitopes of the islet cell autoantigen IA-2 eluted from HLA-DR4"; <i>The Journal of Clinical Investigation</i>; Vol. 104, No. 10; pp. 1449-1457; November 1999; XP-002323392.</p>
	<p>Durinovic-Bello, I., et al; "Predominantly Recognized Proinsulin T Helper Cell Epitopes in Individuals With and Without Islet Cell Autoimmunity"; <i>Journal of Autoimmunity</i>; Vol. 18; pp. 55-66; (2002);</p>

06.28.05